

IMPACT DELIVERY OF REDUCED GREENHOUSE GASES ON EARLY MARS R.M. Haberle¹ and K. Zahnle¹, ¹Space Science and Astrobiology Division, NASA/Ames Research Center, Moffett Field, CA, 94086, Robert.M.Haberle@nasa.gov

Introduction. While there is abundant evidence for flowing liquid water on the ancient Martian surface, a widely accepted greenhouse mechanism for explaining this in the presence of a faint young sun has yet to emerge. Gases such as NH_3 , CO_2 alone, SO_2 , clouds, and CH_4 , have sustainability issues or limited greenhouse power. Recently, Ramirez et al. [1] proposed that CO_2 - H_2 atmospheres, through collision induced absorptions (CIA), could solve the problem if large amounts are present (1.3-4 bars of CO_2 , 50-20% H_2). However, they had to estimate the strength of the H_2 - CO_2 interaction from the measured strength of the H_2 - N_2 interaction. Recent ab initio calculations show that the strength of CO_2 - H_2 CIA is greater than Ramirez et al. assumed [2]. Wordsworth et al. [2] also calculated the absorption coefficients for CO_2 - CH_4 CIA and show that on early Mars a 0.5 bar CO_2 atmosphere with percent levels of H_2 or CH_4 can raise mean annual temperatures by tens of degrees Kelvin. Freezing temperatures can be reached in atmospheres containing 1-2 bars of CO_2 and 2-10% H_2 and CH_4 . The new work demonstrates that less CO_2 and reduced gases are needed than Ramirez et al. originally proposed, which improves prospects for their hypothesis.

If thick weakly reducing atmospheres are the solution to the faint young sun paradox, then plausible mechanisms must be found to generate and sustain the required concentrations of H_2 and CH_4 . Possible sources of reducing gases include volcanic outgassing, serpentinization, and impact delivery; sinks include photolysis, oxidation, and hydrogen escape. The viability of the reduced greenhouse hypothesis depends, therefore, on the strength of these sources and sinks.

Sources. Volcanic outgassing of reduced gases is possible given that the Martian mantle appears to be more reducing than the Earth's [3,4]. Oxygen fugacities in Martian meteorites range from Iron-Wüstite (IW) all the way up to the Quartz-Fayalite-Magnetite (QFM) buffer [3]. If the early Martian mantle was at the low end of this range then a greater fraction of H_2 , CH_4 , and CO would have been included in the outgassed materials. However, the source of CO_2 would then rely more on the oxidation of CO from the photolysis products of water rather than direct outgassing.

Serpentinization is a mechanism in which ultramafic minerals (e.g., olivine) are hydrothermally altered to produce serpentine and magnetite, liberating H_2 in the process. If CO_2 is present in the water it can react with H_2 to produce CH_4 . Thus serpentinization

can produce both H_2 and CH_4 . Serpentine deposits have been identified on the surface [5] and extensive crustal serpentinization may have taken place early in the planets history [6].

Impacts might also be a source of reduced gases. The intense heat and rapid chemistry following an impact could produce H_2 and CH_4 depending on the composition, size, and entry velocity of the impactor, as well as the composition and strength of the target material [7,8,9].

Sinks. Sinks for reduced gases are more easily quantifiable. H_2 escapes while CH_4 is photolyzed and/or oxidized. If H_2 escapes at the diffusion limit, a simple analytical expression can be used to calculate escape rates as a function of mixing ratio and exobase temperature. Sinks for methane can be expressed in terms of lifetimes. Thus, models can be constructed with simple expressions for sources and sinks to estimate the potential for the development of reduced greenhouse gases on early Mars.

A Simple Model. In this work we focus on the production of reduced gases by impacts. Impact production is the least well understood source and the model we construct is meant to assess its potential. A sample production plot for H_2 is shown in Figure 1.




Figure 1. H_2 mixing ratios (%) vs impactor size (km). Dotted horizontal lines labeled T=270, 266, etc. correspond to the global mean temperatures from [2] for a 1 bar CO_2 atmosphere with the indicated hydrogen mixing ratio. Melt is the volume of melt material, and V is the volume of the impactor.

In this figure we have computed H_2 production by assuming a serpentinization-like process that uses the iron in an impactor to make H_2 . We assume a 10% mole fraction of iron, and that one H_2 molecule is produced for every three iron atoms that react with water. We show two curves: one for H_2 generated from the impactor alone ($Melt=1 \times V$, where V is the volume of the impactor), and one where the H_2 comes from a volume target material equal $10 \times V$.

For small melt volumes, significant H_2 production occurs only for sizeable impactors, i.e., impactors greater than ~ 150 km. For larger melt volumes hydrogen production increases sharply for impactors greater than ~ 70 km. Larger impactors occur less frequently than the smaller impactors so the next step is to map our production curves to crater distribution curves and sum up over all impactors to get the total input. We also need to add the sinks discussed above and include a timing profile. We can use the same approach for CH_4 . Our goal is to determine if impacts can build up and sustain the required reduced gas concentrations for solving the faint young sun paradox. Results of this simple model will be presented at the meeting.

References:

[1] Ramirez et al. (2014) *Nature Geo.* 7, 59-63. [2] Wordsworth, et al., (2017), *Geophys. Res. Lett.*, In press. [3] Wadhwa, M. (2008) *Rev. Mineral. and Geochem.* 68,493-510. [4] Hirschmann et al. (2008), *Earth and Planet. Sci. Lett.* 270, 147–155. [5] Ehlmann et al. (2010), *Geophys. Res. Lett.* 37, 6. [6] Chassefière et al. (2013), *J. Geophys. Res.*, 118, 1123-1134. [7] Schaefer and Fegley (2007), *Icarus*, 186, 462–483. [8] Schaefer and Fegley (2010), *Icarus*, 208, 438–448. [9] Hashimoto et al. (2007), *J. Geophys. Res.*, 112, E05010.