



Back in the lab, sediment and source rock samples were prepared for mineralogical and geochemical analysis. Laboratory methods following the field campaign for water and sediment analysis included thin section analysis of source rocks [11], grain size and shape analysis, XRD, XRF, ICP-MS, TEM, and image processing and analysis. The sedimentology of split cores was also logged (Fig. 2), and the analysis of element concentration of cores was performed at regular intervals along the core using a handheld X-ray fluorescence spectrometer.



**Figure 2.** Sediment core from location 2B with the sedimentology and sediment color interpreted on the left panel.

**Results:** In the field, three successful cores were obtained (Fig.1b). At each drill location, groundwater samples were sampled from shallow boreholes that intersected the water table. Offshore surface sediments were additionally collected. The deepest sediment core (~1.5 m) of the 2021 field campaign displays a transition from fluvial to lacustrine sediments (Fig. 2). Importantly, in the lower lacustrine sediments, distinct layers are identified with variations in sediment color.

Preliminary mineralogical analysis of source rocks suggests that the dominant lithology around Lake Sandvatn is basaltic [11]. The XRD results from lake surface sediments display a sediment mineralogy that is dominated by primary basaltic minerals (e.g., plagioclase and pyroxene, Fig. 3) and little to no secondary clay minerals are observed in the bulk sediment. The lacustrine sediments buried at depth display a similar mineral assemblage, but show an increase in X-ray amorphous materials, as identified from an increase in the scatter above the background in the XRD pattern (Fig. 3). There is little evidence for clay minerals in the bulk samples from buried lacustrine sediments.



**Figure 3.** XRD patterns for bulk surface and buried sediments.

**Implications:** Early results from the 2021 DIGMARS field campaign suggest that sediments generated and transported downstream can ultimately carry a mineralogical assemblage consistent with a pristine basaltic progenitor. It is not until these sediments are buried and interacting with groundwater that the mineralogy begins to show evidence for alteration. Early diagenesis in buried sediments from Lake Sandvatn suggests that groundwater discharge is the principal process preserved in the secondary mineralogical record of these sediments. Moreover, aqueous samples show that groundwater discharge and mixing lake waters creates a unique geochemical gradient in the subsurface. This combined with TIR mapping from around the lake illustrate that groundwater discharge in this environment can vary spatially, and almost certainly, temporarily.

When this terrestrial reference frame is applied to regions of Mars (e.g., Gale crater), it becomes clear how early diagenesis from sediments interacting with percolating groundwaters could have a profound impact on interpretations of the sedimentary history of lacustrine environments. Sediments from the ancient lake of Gale crater likely interacted with discharging groundwater and this unique subsurface mixing zone likely played a key role in shaping the sedimentary rock record.

**DIGMARS 2022:** The second field campaign for DIGMARS will take place in the summer of 2022 and will target lakes that are influenced by the discharge of hydrothermally influenced groundwater fluids. A few lakes in southwest Iceland are the target location for this trip, as this region is characterized by both cool and warm groundwater sources. In addition to a second field site, the DIGMARS 2022 drill rig will be deployed from a boat, with the hopes of obtaining deeper sediment cores (e.g., 3 m).

**Acknowledgments:** This work was supported by NASA SSW grant 19-SSW19-0300. We are also grateful to the Iceland Space Agency for their support in the field.

**References:** [1] Grotzinger et al., 2015, *Science* 350(6257), [2] Goudge et al., 2015, *JGR: Planets* 120(4), [3] Mangold et al., 2015, *Science* 374(6568), [4] Meinikmann, Lewandowski, and Nützmänn, 2013, *Journal of Hydrology* 502, [5] Hurowitz et al., 2017, *Science* 356(6341), [6] Rampe et al., 2017, *EPSL* 471, [7] Rampe et al., 2020, *JGR: Planets* 125(12), [8] Bristow et al., 2021, *Science* 373(6551), [9] Thorpe et al., *submitted. JGR: Planets*, [10] Thorpe et al., *JGR: Planets* 126(2), [11] Putnam et al., *this conference*.