

DELIVERY OF ICY PLANETESIMALS TO INNER PLANETS IN THE PROXIMA CENTAURI SYSTEM

S. I. Ipatov, Vernadsky Institute of Geochemistry and Analytical Chemistry (Kosygina st., 19, Moscow 119991, Russia, siipatov@hotmail.com).

The model and initial data used for calculations: The model of migration of planetesimals initially located in the feeding zone of the exoplanet *c* with a semi-major axis $a_c=1.489$ AU in the Proxima Centauri system was studied. The aim of these studies is to compare the delivery of icy planetesimals to potentially habitable planets in the Proxima Centauri system and in our Solar System. Integration of the motion of planetesimals and exoplanets was calculated with the use of the symplectic code from [1] for a star with a mass equal to 0.122 of the solar mass and two exoplanets. It was considered that the exoplanet *b* is located in a habitable zone. In the main series *M* of calculations, based on recent observational data, the following initial semi-major axes, eccentricities, inclinations and masses of two exoplanets were considered: $a_b=0.04857$ AU, $e_b=0.11$, $m_b=1.17m_E$, $a_c=1.489$ AU, $e_c=0.04$, $m_c=7m_E$, $i_b=i_c=0$, where m_E is the mass of the Earth. In the series *F* of calculations, based on older observations, it was considered that $a_b=0.0485$ AU, $a_c=1.489$ AU, $m_b=1.27m_E$, $m_c=12m_E$, $e_b=i_b=0$, $i_c=e_c/2=0.05$ rad or $i_c=e_c=0$. In each calculation variant, initial semi-major axes of orbits of 250 exocomets were in the range from a_{\min} to $a_{\min}+0.1$ AU, with a_{\min} from 1.2 to 1.7 AU with a step of 0.1 AU. Initial eccentricities e_o of orbits of planetesimals equaled to 0.02 or 0.15 for the *M* series, and equaled to 0 or 0.15 for the *F* series of calculations. Initial inclinations of orbits of the planetesimals equaled to $e_o/2$ rad. Considered time interval exceeded 50 Myr. Based on the obtained arrays of orbital elements of migrated planetesimals and exoplanets stored with a step of 100 yr, I calculated the probabilities of collisions of planetesimals with the exoplanets. The probabilities of collisions were calculated also with the unconfirmed exoplanet *d* ($a_d=0.02895$ AU, $m_d=0.29m_E$, $e_d=i_d=0$). The calculations were made similar to those in [2-4].

Probabilities of collisions of planetesimals with the exoplanet *c*: For the *M* series of calculations, the values of the probability p_c of a collision of one planetesimal, initially located near the exoplanet *c*, with this exoplanet were about 0.1-0.3, exclusive for $a_{\min}=1.4$ AU and $e_o=0.02$ when p_c was about 0.6. For the *F* series of calculations at $i_c=e_c=0$ and $e_o=0.15$, p_c was about 0.06-0.1. For $i_c=e_c/2=0.05$ and $e_o=0.15$, p_c was about 0.02-0.04. For both series of calculations, most of planetesimals were usually ejected into hyperbolic orbits in 10 Myr. Usually there was a small growth of p_c after 20 Myr. In some calculations a few planetesimals could still move in elliptical orbits after 100 Myr. The number of planetesimals ejected into hyperbolic orbits was greater by a factor of several than the number of planetesimals collided with exoplanets. Therefore, a cometary cloud similar to the Oort cloud can exist in the Proxima Centauri system.

Probabilities of collisions of planetesimals with the exoplanets *b* and *d*: For the *M* series of calculations, the probability p_b of a collision of one planetesimal, initially located near the orbit of the exoplanet *c*, with the exoplanet *b* was non-zero in 5 among 18 variants at $e_o=0.02$ and in 3 among 6 variants at $e_o=0.15$. At $e_o=0.02$ for the five variants, p_b equaled to 0.004, 0.004, 1.28×10^{-5} , 0.00032 и 9.88×10^{-5} . At $e_o=0.02$ the mean value of p_b for one of 4500 exocomets equaled to 4.7×10^{-4} , but among them there were two planetesimals with $p_b \approx 1$. At $e_o=0.15$ for three variants, p_b equaled to 0.008, 0.004 and 3.6×10^{-6} . The mean value of p_b for one of 1500 exocomets equaled to 2.0×10^{-3} , but among them there were three planetesimals with $p_b \approx 1$. The mean value of the probability p_d of a collision of a planetesimal with the exoplanet *d* equaled to 2.7×10^{-4} and 2.0×10^{-3} at $e_o=0.02$ and $e_o=0.15$, respectively. For the *M* series, the mean values of p_b and p_d averaged over 6000 planetesimals equaled to 8.5×10^{-4} and 7.0×10^{-4} . For all three considered variants of the series *F* at $e_c=0.1$ and $e_o=0.15$, the values of p_b were in the range 0.008-0.019. For other calculations of the *F* series, $p_b=0$. Only one of several hundreds of planetesimals reached the orbits of the exoplanet *b* and *d*, but the probabilities p_b and p_d of a collision of one planetesimal with these exoplanets (averaged over thousands planetesimals) are greater than the probability of a collision with the Earth of a planetesimal from the zone of the giant planets in the Solar System. The latter probability for most calculations with 250 planetesimals was less than 10^{-5} per one planetesimal [5]. Therefore, a lot of icy material could be delivered to the exoplanets *b* and *d*.

Acknowledgments: For studies of formation of exoplanets and of the ejection of exocomets into hyperbolic orbits, the author acknowledges the support of Ministry of Science and Higher Education of the Russian Federation under the grant 075-15-2020-780 (N13.1902.21.0039). Migration of icy planetesimals to exoplanets located in the habitable zone was carried out as a part of the state assignments of the Vernadsky Institute of RAS № 0137-2021-0004.

References: [1] Levison H.F., Duncan M.J. (1994) *Icarus* 108:18-36. [2] Ipatov S.I., Mather J.C. (2004) *Annals of the NYAS* 1017:46-65. <http://arXiv.org/format/astro-ph/0308448>. [3] Ipatov S.I., Mather J.C. (2004) *Advances in Space Research* 33: 1524-1533. <http://arXiv.org/format/astro-ph/0212177>. [4] Ipatov S.I. (2019) *Solar System Research* 53: 332-361. <http://arxiv.org/abs/2003.11301>. [5] Ipatov S.I. (2020) *EPSC2020-71*, <https://doi.org/10.5194/epsc2020-71>.