

Regional Geologic Mapping of the Oxia Planum Landing Site for ExoMars.

S. Adeli¹, A. Nass¹, D. Tirsch¹, S. Acktorries¹, S. Steffens¹, and E. Hauber¹, ¹Institute of Planetary Research, German Aerospace Center (DLR), Rutherfordstr. 2, 12489, Berlin, Germany, Solmaz.Adeli@dlr.de.

ABSTRACT

The ExoMars rover mission will deploy a stationary surface platform and a rover in Oxia Planum (OP), a region at the transition between the cratered highlands of Mars and the ancient and filled impact basin, Chryse Planitia. While the fundamental geologic characteristics of the area have been investigated during the landing site selection process, detailed geologic or morphostratigraphic mapping is still missing. To fill this knowledge gap, two complementary mapping approaches were initiated by the ExoMars RSOWG: (1) Local HiRISE-scale mapping of the landing ellipse(s) area [1]. (2) Regional mapping at CTX-scale [2 & this study] which will provide a more synoptic view of the wider landing site within OP, enabling the contextualization of the units within the stratigraphy of western Arabia Terra and Chryse Planitia. It is also expected that this map will serve as a geologic reference throughout the mission and subsequent data analysis.

The study area is located between 16.5°N and 19.5°N, and 22°W to 26°W. The data sets used for mapping include HRSC, THEMIS IR (day and night), CTX, and CaSSIS. Mapping scale in a GIS environment is 1:100,000, which will result in a final printable map at a scale of 1:1M. Mapping started in mid-October 2019. Overall, the identified map units are very similar to those described by [3]: The spatially most widespread units are the phyllosilicate-bearing unit that is the prime ExoMars target (with distinctly enhanced THEMIS nighttime temperatures compared to its surroundings) (also described in [4]), a dark resistant unit of possibly volcanic or sedimentary origin, and a mantling unit that was likely emplaced by eolian processes. Multiple channels of various morphology and degradation state as well as fan-shaped deposits (with low nighttime temperatures) imply a diverse and possibly long-lived history of surface runoff, perhaps accompanied or replaced by groundwater processes such as sapping. Inverted landforms (channels, impact craters) are the result of intense erosion. Additional mapped features include wrinkle ridges and lobate scarps (delineating a basin-like depression in the central mapping area), remnant buttes in the northwestern portion of the mapping area (i.e. towards Chryse Planitia), craters and their ejecta blankets, and fields of eolian bedforms and secondary craters. At the time of writing, the mapping is incomplete and only initial and limited conclusions can be drawn. Overall, the mapping confirms previous geologic analyses. However, some features (e.g., contractional structures, channels, possible sapping landforms) need further attention as they may provide important constraints on the tectonic and aqueous evolution of the ExoMars landing area.

References: [1] Sefton-Nash et al. *LPSC* (2020). [2] Hauber et al., *LPSC* (2020) [3] Quantin et al. *Astrobiology* (under review). [4] Mandon et al., *Astrobiology* (under review).

Acknowledgments: The data used in this study are available via the Planetary Data System (PDS) of NASA, the Planetary Science Archive (PSA) of ESA, and additional data repositories at DLR and the USGS.




Figure Caption: Geologic map of the ExoMars rover landing site in Oxia Planum, at CTX-scale. Distance between two longitudes is ~55.75 km.