

LUNAR CRATER MATURITY ANALYSIS IN PYTHON: DEVELOPING A TOOLKIT FOR EJECTA ANALYSIS. K.

A. Carr¹, O. Azubuike¹, A. Tran², C. Carreira², C. Alfaro¹, B. Greenhagen³, G.W. Patterson³, A.M. Stickle³, P. Prem³, J. T. S. Cahill³, C. J. Tai Udovicic⁴. ¹University of Maryland, Baltimore County, Baltimore, MD (kat50@umbc.edu). ²Johns Hopkins University, Baltimore, MD. ³Johns Hopkins University Applied Physics Laboratory, Laurel, MD (angela.stickle@jhuapl.edu). ⁴Northern Arizona University, Flagstaff, AZ.

Introduction: The lunar surface is covered by impact craters, which were formed throughout solar system history. The expression of craters can change with age and affect what is seen in remote sensing data. Examining a variety of impact craters can provide information about how the ejecta of these craters evolves with exposure to the space environment. Ultimately, the goal of this research is to be able to better understand the evolution of the lunar surface by understanding the evolution of impact ejecta. Here, we describe a toolkit developed for the purpose of being able to evaluate the ejecta of young craters on the Moon.

Background: *Maturity of lunar soils.* Maturity in this context refers to a qualitative estimate of the length of time a specific region of interest has experienced surface exposure. By looking at the maturity of craters and their ejecta, important questions such as when these impacts occurred, and what the surface around the crater is like, can be answered. Changes in the physical properties of lunar soil with age are usually quantified in terms of specific indices (e.g., Optical Maturity (OMAT) [1]). Evaluating maturity indices at different wavelengths [e.g., 1-8] provides a more complete understanding of how ejecta degrade on the lunar surface. Crater ejecta can reveal a lot of information about how a given crater formed, and can more broadly tell us more about the composition of the Moon and how the surface evolves with time.

Methods of evaluating maturity. Current available tools that are used to evaluate crater maturity include, ISIS, MATLAB, and IDL. To varying degrees, these tools have been utilized with a wide variety of datasets to examine maturity of lunar soils (e.g., LROC WAC [2], Diviner [3], M3 [4], Mini-RF [5,6], and OMAT [1]). As a result, the workflow involved with conducting this type of analysis can be complex, requiring significant manual intervention. The goal of the Lunalyze toolkit is to streamline workflow for this type of analysis, primarily through the automation of tasks.

Method: The Lunalyze toolkit uses Python to allow users to be able to open and run select scripts. Python was chosen because of its versatility as a computing language, the fact that it is open source, and that it is relatively easy to access and learn how to use. Lunalyze builds on existing Python packages [e.g., 9], but includes new features in Jupyter Notebooks, which allow the user to get real time feedback on which segments of the scripts are working and which ones are not. We also included virtual anaconda environments for Windows, Linux, and MacOS. Currently, Lunalyze can read in a variety of image file types and convert them to other file types as needed. The image conversion is done using a graphical user

interface (GUI) that removes the step of having someone go and edit the source code (Figure 1).




Figure 1: An image of the GUI for image conversion.

A variety of different data types can be converted as needed, including standard PDS formats, image cubes and MATLAB files.

Current development efforts for Lunalyze analysis capabilities include crater detection, rim and floor detection, and shape file generation. This work builds on the open-source Python package Craterpy [9]. The Lunalyze interface allows a user to look at regions of interest based on specified coordinates and perform analysis for a given dataset. This analysis includes options for looking at the crater floor, rim, and ejecta individually. Ongoing development will facilitate the analysis of craters (and their ejecta) across a variety of lunar data sets.

The current use-case for Lunalyze is to provide a user-friendly and efficient means of evaluating the maturity of Copernican crater ejecta. Our objectives are twofold: 1. To evaluate whether maturity indices at different wavelengths can be correlated, and 2. To provide a new, open-source toolkit to the community that can be used to study lunar craters across a variety of data sets. Stickle et al. [5] showed that the maturity of ejecta blankets for lunar craters could be tracked across wavelengths for craters of varying ages. That initial work used radial profiles that were placed and calculated manually to evaluate maturity indices in LRO and Kaguya data. To increase the time and efficiency of conducting such analyses, Lunalyze provides a user-friendly interface that includes automation options.

Like the GUI image converter, we have been working on ways to allow users to mask craters without having to edit the source code. This feature has been developed by using two scripts that work back-to-back with each other. The first script loads the image that the user would like to place a circular mask over, and the user can click on the center of the crater floor and the crater rim. This script records the coordinates of where the user has clicked on the picture, and then these values are transferred over to the second script, which creates a circular mask based off of the clicks that the user has provided. It centers the circle on the first set of coordinates that the user has provided, and then uses the second click to get the radius of the circle that needs to be created for the mask. By using this option, we are able to begin looking at other kinds of data covering crater ejecta blankets, including, for example, the Diviner rock abundance (e.g., Figure 5).

Discussion and Future Work: The development of Lunalyze is ongoing and is currently focused on options related to detecting crater rims and extracting radial profiles from selected regions of crater ejecta blankets. This capability is being tested on select lunar craters. An example is the crater Dufay B (Figure 2). Dufay B was chosen as a stressing case due to its irregular rim and non-circular shape. These factors can pose a significant challenge for an automated tool. The option to manually draw the crater rim is in development. However, automated options that can address this case are being considered and evaluated.

Lunalyze has been designed to accommodate analysis across multiple relevant datasets (e.g., Diviner, LOLA, and Mini-RF). While current tested capability for viewing and analyzing craters is restricted to image data (e.g., Figure 2) we are slowly creating graphical models. Using a 750nm Clementine global mosaic we can now analyze the visible albedo value of craters (Figure 4). Diviner data allows examination of rock abundance and nighttime soil temperature. We hope to create many more of these models by utilizing the data provided by various database on the craters we wish to study in the near future. Multi-band datasets still pose a challenge for integration in our analysis framework and is another focus of ours.

Acknowledgments: This work was supported by APL's CIRCUIT leadership, research, and mentoring program, and by the NASA Lunar Data Analysis Program, grant number 80NSSC19K0370, the Mini-RF radar project, and the LRO Diviner Lunar Radiometer project.

References: [1] Grier et al. (2001) *J. Geophys. Res.* 106(E12), 32847-32862; [2] Denevi et al. (2014) *J. Geophys. Res.: Planets* 119(5) 976-997; [3] Lucey et al. (2017) *Icarus*, 283, 343-251; [4] Nettles et al. (2011) *J. Geophys. Res.: Planets* 116(E9); [5] Stickle, A.M. et al. (2016) LEAG Abstract #5055; [6] Neish et al. (2013) *J. Geophys. Res.: Planets*, 118, 2247-2261; [7] Lucey, P.G., et al. (2000) *J. Geophys. Res.*, 105, 20,377-20,386; [8] Greenhagen, B. T., et al. (2010) *Science*, 329, 1507-1509; [9] Tai Udovicic, C.J.,

(2017) *cjtu/craterpy: craterpy-reactoring* (Version 0.2), Zenodo.




Figure 2: A 750nm Clementine global mosaic of the highlands crater Dufay B, one of the craters of interest in this study Dufay B (19.8 km, 8.3°N, 171°E) is categorized as “intermediate” in age [2].




Figure 3: Dufay B with a circle estimating the crater overlain. The white circle represents the automated determination of the crater planform in Lunalyze, showcasing still needed adjustments.




Figure 4: A histogram of the albedo values of Dufay B's ejecta. The X-axis represent the albedo values and Y-axis is the pixels.




Figure 5: A histogram of Rock Abundance in the ejecta of Dufay B, showing that the ejecta is not particularly rocky (low value on the X-axis).