

**COLD ORIGINS: PREBIOTIC CHEMISTRY AT THE ICE MATRIX.** C. Menor-Salván<sup>1,2</sup> and M. Roig Marín-Yaseli<sup>2</sup>, <sup>1</sup>Department of Chemistry and Biochemistry, Georgia Institute of Technology, Atlanta, GA, 30332-0400, USA, <sup>2</sup>Centro de Astrobiología (CSIC-INTA), Ctra. Torrejón-Ajalvir km 4, Torrejón de Ardoz, 28850, Spain.

**Introduction:** The origin of nucleobases and other heterocycles is a classic question in the chemistry of the origins of life. The construction of laboratory models for the abiotic synthesis of nitrogen heterocycles in plausible natural conditions also aids the understanding and prediction of chemical species in the Solar System. If we support the idea of a prebiotic origin of nucleobases, a “cold origin of life” [1] or “ice world” is the preferred prebiotic environment because the nucleobases would be stable under cold conditions, and such an environment supports the relevance of eutectic water solutions of reactants in an ice matrix. During the last years, we performed a series of experiments [2,3] which demonstrate the feasibility of the prebiotic synthesis of nucleobases at the ice matrix in the range of existence of eutectic solutions of organic solutes.

**Results:** The irradiation (by spark discharges or UV irradiation) of an atmosphere containing methane/ nitrogen or acetylene, lead to the formation of organic precursors whose fate depends on the environmental conditions.




Figure 1. Origin of nitrogen bases in urea water-ice solution under methane atmosphere.

If a water pool is present at a temperature sufficiently low as freeze-thaw cycles are set, products of atmospheric irradiation could react with organic solutes, especially in the concentrated brines formed during the freezing process. The low temperature and freezing gives different results from the classic prebiotic chemistry simulations. Thus, the irradiation of a system composed by methane-rich atmosphere and an ice pool, lead to the concentration of polycyclic aromatic hydrocarbons in water. If a nitrogen solute is present in eutectic solutions, as urea, the formation of pyrimidines (mainly uracil) and purines (mainly adenine) (Figure 1).

If acetylene (a molecule that has received little attention in the development of Prebiotic Chemistry) is present in the atmosphere, its photolysis products could

react with efficient condensing agents, as urea, leading to the efficient formation of nitrogen heterocycles (Figure 2).




Figure 2: Acetylene as precursor of nucleobases.

The experiments performed demonstrated that the synthesis of aromatic hydrocarbons, purines and pyrimidines and other nitrogen heterocycles of potential prebiotic interest (such as triazines and hydantoins) are favoured in the ice matrix by classic cyanide and cyanoacetylene pathways or by condensation of acetylene photolysis products. Despite these results, the experimental prebiotic chemistry in the solute-concentrated solutions that fill the space confined by ice matrix has received relatively little attention in the elaboration of the models for the origin of organics in Solar System bodies and prebiotic evolution. Consequently, it is necessary to perform more experiments under plausible prebiotic conditions, especially if geochemical models support stable icy environments on the prebiotic Earth. The *ice world* constitutes an interesting prebiotic chemistry scenario that awaits further investigation.

#### References:

- [1] Chyba, C.F., and Phillips, C.B. (2002) *Orig. Life Evol. Biosph.*, 32, 47-67.
- [2] Tobie, G., Grasset, O., Lunine, J.I., Mocquet, A., and Sotin C. (2005) *Icarus*, 175, 496-502.
- [3] Moulton, V., Gardner, P.P., Pointon, R.F., Creamer, L.K., Jameson, G.B., and Penny, D. (2000) *J. Molec.Evol.*, 51, 416-421.
- [4] Miller, S.L., and Orgel, L. *The Origins of Life on the Earth*, Prentice Hall, New Jersey, 1974.
- [5] Bada, J.L. (2004), *Earth Planet. Sci. Lett.*, 226, 1-15.
- [6] Menor-Salván, C., Ruiz-Bermejo, M., Guzmán, M.I., Osuna-Esteban, S. and Veintemillas-Verdaguer, S. (2009) *Chem. Eur. J.*, 15, 4411-4418.
- [7] Menor-Salván, C. and Roig Marín-Yaseli, M. (2013) *Chem. Eur. J.*, 19, 6488-6497.