

**MAIN-BELT INFRARED SPECTRAL ANALOGUES FOR (101955) BENNU: GAUSSIAN FITTING TO AKARI SPECTRA OF BENNU-LIKE ASTEROIDS.** L. F. Lim<sup>1</sup>, H. H. Kaplan<sup>2</sup>, V. E. Hamilton<sup>2</sup>, P. R. Christensen<sup>3</sup>, A. A. Simon<sup>1</sup>, D. C. Reuter<sup>1</sup>, J. P. Emery<sup>4</sup>, B. Rozitis<sup>5</sup>, M. A. Barucci<sup>6</sup>, A. Praet<sup>6</sup>, H. Campins<sup>7</sup>, B. E. Clark<sup>8</sup>, M. Delbo<sup>9</sup>, J. Licandro<sup>10</sup>, R. D. Hanna<sup>11</sup>, S.A. Sandford<sup>12</sup>, E.S. Howell<sup>13</sup>, D. S. Lauretta<sup>13</sup>, <sup>1</sup>Goddard Space Flight Center, Greenbelt, MD, USA (lucy.f.lim@nasa.gov), <sup>2</sup>Southwest Research Institute, Boulder, CO, USA, <sup>3</sup>Arizona State University, Tempe, AZ, USA, <sup>4</sup>University of Tennessee, Knoxville, TN, USA, <sup>5</sup>Open University, Milton Keynes, UK, <sup>6</sup>LESIA, Paris Observatory Meudon, France, <sup>7</sup>University of Central Florida, Orlando, FL, USA, <sup>8</sup>Ithaca College, Ithaca, NY, USA, <sup>9</sup>CNRS, France, <sup>10</sup>Instituto de Astrofísica de Canarias, Tenerife, Spain, <sup>11</sup>University of Texas, Austin, TX, USA, <sup>12</sup>Ames Research Center, Mountain View, CA, USA. <sup>13</sup>Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ, USA.

**Introduction:** The Origins, Spectral Interpretation, Resource Identification, and Security–Regolith Explorer (OSIRIS-REx) mission has measured the spectrum of asteroid (101955) Bennu in reflectance (OVIRS instrument; [1]) and thermal emission (OVIRS and OTEs instruments; [2]). Here we place the global average spectrum of Bennu [3] in the context of the wider asteroid population as represented by infrared reflectance spectra from the AKARI mission [4].




Figure 1: Dynamical context of AKARI and selected Spitzer asteroids in the main belt relative to the probable Bennu source region (orange ellipse). Asteroids with Bennu-like spectral shapes in the 2.6–3.5  $\mu\text{m}$  region can be found at a wide range of semimajor axes and inclinations.

On dynamical grounds (101955) Bennu has been considered most likely to have originated in the inner main belt families of (495) Eulalia (C-type, semimajor axis  $a = 2.49$  AU) or (142) Polana (B-type,  $a = 2.42$  AU) [6] (Fig. 1). However, neither Eulalia, nor Polana, nor their family members were observed spectroscopically either with AKARI or with the Spitzer IRS.

**B-type main belt asteroids in the AKARI spectral catalogue:** (2) Pallas, (704) Interamnia, and (24) Themis were observed spectroscopically by AKARI. Although all three asteroids are dynamically distant from the Polana/Eulalia complex and the  $\nu_6$  secular resonance, Pallas and Interamnia are relatively close spectral matches in the 2.6–3.5  $\mu\text{m}$  wavelength region, in which Bennu's strongest spectral feature is located (Figs. 2 and 3) [3].




Fig. 2: OVIRS spectrum of Bennu vs. AKARI spectra of B-type asteroid (2) Pallas, B- or Cb-type asteroid (704) Interamnia, and inner-main-belt Cgh-type asteroid (51) Nemausa




Figure 3: Preliminary four-Gaussian fits to the AKARI spectra of (2) Pallas and (704) Interamnia. The Gaussian at  $\sim 2.9$   $\mu\text{m}$  is much stronger in the spectrum of Pallas as a fraction of the total band area. Gaussian deconvolutions following Hiroi et al. 2018, 2019; Kaplan et al. 2019 [7,8]

Like Bennu, Pallas and Interamnia both have band minima at  $\lambda_{2.7} = 2.74 \pm 0.01$   $\mu\text{m}$  [3, 4]. The shape of

Bennu's 2.7- $\mu\text{m}$  band is a substantially better match to that of Pallas in the 2.85–3.0  $\mu\text{m}$  region. Preliminary Gaussian fits (Fig. 3) show that the difference in shape can be explained by the size of the Gaussian at  $\sim 2.89$   $\mu\text{m}$ . See Praet et al. (this meeting, abstract #1058) for Gaussian fits to the OVIRS spectrum of Bennu [9].

In contrast, B-type (24) Themis is a comparatively poor match to Bennu in this region and also contains a deep 3.1- $\mu\text{m}$  band [10,11] not matched by corresponding structure in Bennu's spectrum.

**Other C-complex and Xk-type Main Belt analogues to Bennu:** Several other asteroids in the AKARI catalog are similarly close spectral matches to Bennu based on a combination of  $\chi^2$  and correlation tests. (476) Hedwig is notable for the similarity of its spectral shape (Figs. 4, 5) in spite of its VNIR classification as an Xk or P-type asteroid rather than a member of the C-complex.




Figure 4: Bennu-like band shapes in Xk-type asteroid 476 Hedwig and Ch-type outer-main-belt asteroid 49 Pales

We note that as with (24) Themis, there are also many C-type objects that are very unlike Bennu in the 2.7- $\mu\text{m}$  region, such as (94) Aurora, (52) Europa, and (451) Patientia. The Xc-type (21) Lutetia, despite being a member of the low-inclination inner-main-belt population, is also not an analogue to Bennu in this spectral region.

**Summary and Conclusions:** Spectral analogues to Bennu in the 2.6–3.2  $\mu\text{m}$  region are widespread among the large main-belt asteroids in the AKARI catalogue and are commonly dynamically distant from the most likely Bennu source regions in the main belt. Pallas is a close spectral analogue to Bennu in spite of its high inclination. At the AKARI spectral resolution, the Bennu-like spectral shape can be represented by four




Figure 5: Four-Gaussian fit to the AKARI spectrum of Xk-type asteroid 476 Hedwig. As with the fit to (2) Pallas, the Gaussian close to 2.9  $\mu\text{m}$  represents nearly 30% of the total band area.

Gaussians, one of which falls close to 2.89  $\mu\text{m}$  and represents a substantial fraction of the total band area.

Further observations, likely space-based, will be needed in order to determine whether this spectral shape is prevalent among the main-belt populations with the most straightforward dynamical pathways to Bennu-like orbits: the low-inclination inner-main-belt C-complex asteroid families, or the low-inclination inner-main-belt population more generally.

**Acknowledgments:** This material is based upon work supported by NASA under Contract NNM10AA11C issued through the New Frontiers Program. We are grateful to the entire OSIRIS-REx team for making the encounter with Bennu possible.

**References:** [1] Reuter, D. C. et al. (2018) *Space Science Reviews*, 214, 54. [2] Christensen, P. R. et al. (2018) *Space Science Reviews*, 214, 87. [3] Hamilton, V. E. et al. (2019) *Nature Astronomy*, 3, 332-340. [4] Usui, F. et al. (2019) *Publ. Astron. Soc. Japan* 71, 1 (1–41). [5] Houck, J. R., et al. (2004) *ApJS* 154, 18. [6] Bottke, W. F., et al. (2015) *Icarus* 247, 191. [7] Hiroi, T. et al. (2019) *LPSC 50*, #1129. [8] Kaplan, H. H., et al. (2019) *M&PS*, 54: 1051-1068. [9] Praet, A. et al. (2020) *LPSC 51*, #1058. [10] Rivkin, A.S. and Emery, J.P. (2010) *Nature* 464, 1322. [11] Campins, H. et al. (2010) *Nature* 464, 1320.