

AN ATTEMPT TO DETERMINE MINIMUM AREAS FOR USEFUL MODEL AGES. D. Hill¹, O. Fox, A. Hager¹, O. Ukiwo¹, A. Duffy¹, C. Akers¹ and A. D. Schedl¹ (Department of Physics, West Virginia State University, schedlad@wvstateu.edu.)

Introduction: We have determined model isochron ages of 44 landslides in Valles Marineris (VM) that are $\geq 400 \text{ km}^2$ in area [1]. Almost 50% of these landslides have age's $\leq 1.0 \text{ Ga}$. If these ages are valid, then it has important implications for the origins of these landslides, i. e., they could not have been initiated by meteorite impact.

Another possibility is that the landslide model ages are unreliable because of resurfacing. Our previous work suggests that landslides composed of chaotic material with undulating topography are susceptible to rapid degradation of impact craters resulting in model ages that are too young, even though the landslide is $\approx 3600 \text{ km}^2$ in area. A critical factor in the reliability of isochron model ages is the area of the feature being dated [6]. This work suggest that reproducible isochron can be determined for areas of $1,000 \text{ km}^2$. The long-term aim of this study is to try to establish a minimum area for useful isochron ages.

Methods: Robbins et al. [7] developed a classification scheme based on the degradation state of impact craters and a database of 400,000 craters. In previous studies, we found that crater degradation states of craters near VM are related to age [8]. 90% of craters of degradations states 1 and 2 are older than 3.5 Ga, whereas degradation state 3 craters range from 1.0 to 3.6 Ga. The number of craters of degradation state 4 near VM are small in size, $\leq 30 \text{ km}$ diameter, and relatively few in number. Preliminary ages degradation state 4 craters are $< 1.0 \text{ Ga}$.

These results suggests that degradation state 3 craters are most likely to be linked with landslides in VM. Our control group were 38 impact craters of degradation state 3, $> 40 \text{ km}$ in diameter, and lying within 2,000 km of the nearest landslide in VM and are drawn from the above database [7]. Only one of these craters was misclassified. Thus, we determined model isochron ages using Craterstats II [9] of 37 craters and crater counted using JMars. Our experimental groups are 40 craters of degradation state 3 lying within 1200 km of landslides in VM, and 10 to 15 km complex craters and 15 to 20 km in diameter. An additional experimental group are degradation state 3 craters 23-40 km in diameter lying within $\pm 30^\circ \text{ N}$ and S latitude lying near Valles Marineris. These craters were selected using the same database, out 20% of the craters that were misclassified. We noted that about 35% of craters contained impacts $\geq 1 \text{ km}$, so we also dated craters using Neukum [10]. We then tested whether our control and experimental group were drawn from the same distribu-

tion using a Kolmogorov-Smirnov statistical test. Our plan is to date additional size ranges of impact craters and additional degradation state 3 craters we have identified $\geq 40 \text{ km}$ diameter near VM. Our choice of studying craters near VM to understand landslide ages in VM are 1) the crater fill and landslide material should be similar in terms of source rocks and material characteristics. 2) The landslides and craters are at nearly the same latitudes so they should have similar climate histories.

Results: Figure 1 shows the age frequency distribution for degradation state 3 craters $\geq 40 \text{ km}$ in diameter. The figure shows an exponential decay in the number of craters over time. Figure 2 shows the Isochron ages of craters 15 to 20 km in diameter. Most of the crater ages are $\leq 1.6 \text{ Ga}$. In Figure 3 Neukum ages [10] were determined using the areal density of craters $\geq 1 \text{ km}$ in diameter which were then substituted for the isochron ages. The curve is considerably more bimodal than figure 2. Figure 4 shows a typical isochron for which a Neukum age was also calculated. The isochron age of 1.5 ± 0.6 was used in Figure 2 and a Neukum age of 3.6 Ga was used for Figure 3. Figure 5 shows the Isochron ages for complex craters 10 to 15 km diameter of degradation state 3 and Figure 6 shows the isochron ages for craters 23 to 40 km in diameter of degradation state 3.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

Interpretation: An exponential decrease in the rate of impact cratering (Figure 1) for Mars is consistent with data from the Earth [2] and the Moon [3]. Thus, the figure 1 is representative of the age-frequency distribution of degradation state 3 craters near VM. The Kolmogorov-Smirnov $D = 0.6189$ for a comparison of the curves in Figure 1 and Figure 2 and Kolmogorov-Smirnov $D = 0.5189$ for a comparison of the curves in Figure 1 and Figure 3. Comparisons of Figure 1 with 4 & 6 are respectively $D = 0.5398$ and $D = 0.3759$. Thus, the probability that the data shown in figures 2, 3 4 & 6 are drawn from the same distribution as the data in figure 1, $P \ll 0.001$. The difference in the distributions is probably the result of erosion. However, the lower D statistic for the curve in figure 3 and the percentage of ages between 3.2-3.6 Ga being nearly the same for figure 1 and 3, suggests that absent the effects of resurfacing the 15-20 km ages and the >40 km ages may be drawn from the same age distribution. Looking at age distributions for different size ranges may resolve the issue of being drawn from the same distribution and a minimum area for crater counting.

References:

[1] A. Hager and A. D. Schedl (2017) *LPS, XLVIII*, Abstract #2076.. [2] D. R. Lowe and Byerly. (2018) *New Astro. Rev.*, 81, 39.. [3] W. K. Hartman (2019) *Geosciences*, 9, 285. [4] Quantin, C. et al. (2004) *Icarus*, 172, 555-572. [5] A. Hager, A. D. Schedl and O. Ukiwo, (2018) *LPS, XLIX*, Abstract #2062. [6] Warner, N. H. et al. (2015) *Icarus*, 245, 198-240. [7] Robbins S. J. et al., (2012) *JGR*, 117, E05004. [8] Duffy, A. and Schedl, A. D. (2015) *LPSC 46th*, Abstract # 2501. [9] Michael, G. G. and Neukum, G. (2010) *EPSL*, 294, 223-229. [10] G. Neukum (1983), *Ph. D. Dissertation, Univ. of Munich*, 186.