

NOBLE GASES IN THE HAMLET METEORITE (LL4).

Sachiko Amari¹, Yukie Sabe², Tomokazu Shiraishi² and Jun-ichi Matsuda². ¹McDonnell Center for the Space Sciences and the Physics Department, Washington University, St. Louis, MO 63130, USA. E-mail: sa@wuphys.wustl.edu. ²Department of Earth and Space Science, Osaka University, Osaka, 560-0043, Japan.

Introduction: Heavy noble gases in primitive meteorites are contained in a very small portion of meteorites. This phase, Q for quintessence, is most likely carbonaceous [1, 2]. However, the exact nature of Q remains enigmatic. In continued effort to better understand Q [3-6], we carried out noble gas studies of the Hamlet meteorite (LL4), both for a bulk sample and an HF-HCl residue.

Experimental: The total of 3.47 g of the meteorite was processed at Washington University in St. Louis, USA. It was alternately treated with HF-HCl and HCl to dissolve silicates. Elemental sulfur was removed with CS₂. The residue comprised 1.11% of the starting material. Noble gases were analyzed using the VG5400 at Osaka University, Japan. The temperatures for step-wise heating were 600, 800, 1000, 1200, 1400, and 1600°C for both samples.

Results and Discussion: The ⁸⁴Kr and ¹³²Xe concentrations in the bulk sample are 1.24×10^{-9} and 1.40×10^{-9} cm³STP/g, respectively. They are 2.2 and 2.8 times higher than those of the same meteorite by Alaerts et al. [7]. Helium-4, ²²Ne and ³⁶Ar concentrations are comparable in the two studies, indicating Q is more abundant in our bulk sample.

The ¹³²Xe concentration of the HF-HCl residue is 9.24×10^{-8} cm³STP/g, indicating that Q was enriched 66 times from the bulk sample. The mass balance calculation indicates that 73 percent of the Xe in the bulk sample remains in the HF-HCl residue. The Xe isotopic composition of the HF-HCl residue indicates that the Xe is essentially Xe-Q and that no diamond was in the residue.

The Ne in the bulk sample is dominated by cosmogenic Ne: the ²⁰Ne/²²Ne and ²¹Ne/²²Ne ratios of the total are 0.812 ± 0.004 and 0.866 ± 0.003 , respectively. The residue, although all silicates were removed, still contains a large amount of cosmogenic Ne (2.467 ± 0.017 and 0.6576 ± 0.006 , respectively), indicating that it contains a significant amount of oxides. In order to determine the ²⁰Ne/²²Ne ratio of Hamlet Ne-Q, we assumed 0.029 as the ²¹Ne/²²Ne ratio of Hamlet Ne-Q because ²¹Ne/²²Ne ratios of Ne-Q from many meteorites are known to be ~ 0.029 [4, 8]. The ²⁰Ne/²²Ne of Hamlet Ne-Q was determined to be 11.0 ± 0.5 . ²⁰Ne/²²Ne ratios of Ne-Q vary in different meteorites: they can be divided into two groups [8, 9]. The first group has ²⁰Ne/²²Ne ratios of 10.11 ± 0.04 , including Ne-Q from Lancé (CO3.5), and Cold Bokkeveld (CM2), and the second group has those of 10.57 ± 0.19 , including Ne-Q from Allende (CV3), Chainpur (LL3.4), Grosnaja (CV3) and Murchison (CM2) [8]. Hamlet Ne-Q obviously falls into the second group.

References: [1] Reynolds J. H. et al. 1978. *Geochim. Cosmochim. Acta*, 42:1775-1797. [2] Ott U. et al. 1981. *Geochim. Cosmochim. Acta*, 45:1751-1788. [3] Matsuda J. et al. 2010. *Geochem. Cosmochim. Acta*, 74:5398-5409. [4] Matsuda J. et al. 2010. *Meteorit. Planet. Sci.*, 45:361-372. [5] Matsuda J. et al. 2010. *Meteorit. Planet. Sci.*, 45:1191-1205. [6] Amari S. et al. 2013. *Astrophys. J.*, 778:37 (39pp). [7] Alaerts L. et al. 1979. *Geochim. Cosmochim. Acta*, 43:1399-1415. [8] Busemann H. et al. 2000. *Meteorit. Planet. Sci.*, 35:949-973. [9] Huss G. R. et al. 1996. *Geochim. Cosmochim. Acta*, 60:3311-3340.