

I-XE SYSTEMATICS OF BRACHINITE-LIKE ULTRAMAFIC ACHONDRITE NORTHWEST AFRICA 5400.

O. Pravdivtseva¹, A. Meshik¹, C. M. Hohenberg¹ and A. J. Irving². ¹Physics Department, Laboratory for Space Sciences, Washington University in St. Louis, St. Louis MO 63130, USA E-mail: oa@physics.wustl.edu, ²Dept. of Earth & Space Sciences, University of Washington, Seattle, WA 98195, USA.

Introduction: The ungrouped brachinite-like achondrite NWA 5400 is a relatively oxidized, metal-bearing ultramafic stone with terrestrial O and Cr isotope composition [1-4]. A comprehensive petrographic, elemental and multi-isotopic study of NWA 5400 and paired NWA 5363 suggested that the mineral assemblage in these meteorites represents a restite after partial melting and extraction of a small amount of silicate from a fertile source rock within ~ 1.5 Myr of the start of the solar system [5]. U-Pb systematics suggested very early differentiation, consistent with evolution of some parent bodies of iron meteorites [6]. Mn-Cr systematics indicated that ⁵³Mn fully decayed at the time of isotopic closure [7]. Here we report I-Xe data for mineral phases separated from NWA 5400.

Experimental: In situ laser analyses on a polished section of NWA 5400 indicated the presence of multiple ¹²⁹Xe rich mineral phases potentially suited for I-Xe dating, consistent with previous results for the whole rock [8]. To prepare mineral separates the meteorite was gently crushed, sieved into grain sizes between 37 and 74 μm and magnetically separated using a Frantz Isodynamic Magnetic Separator at progressively higher current settings. Troilite grains were manually removed from separates after each run. The final fraction, non-magnetic at 1.8A, consisted of mixture of forsteritic olivine and apatite. Aliquots of all samples were saved for mineralogical studies. The resulting 8 samples and absolute age standard Shallowater were sealed under vacuum in quartz tubes and irradiated with thermal neutrons to convert ¹²⁷I into ¹²⁸Xe, receiving ~ 2×10¹⁹n/cm². Samples were wrapped in Pt, the xenon was extracted by stepwise heating in a low blank W-coil and analyzed by high-transmission mass spectrometry. Hot blanks were measured at temperatures ~ 100 °C higher than melting of Pt and were consistently atmospheric in composition.

Results: Two separates were analyzed so far. Olivine/apatite yielded an apparent high-temperature isochron corresponding to closure of I-Xe system 6.6 ± 0.5 Ma before Shallowater, and to an absolute age of 4568.9 ± 0.6 Ma (assuming a Shallowater age of 4562.3 ± 0.4 Ma [8]), within the Pb-Pb age of CAI (4568.2 ± 0.2 Ma [9]). This age may reflect the time of partial differentiation suggested by [5]. The I-Xe system in troilite closed ~ 8 Ma later. Contrary to what was observed for Mn-Cr, the I-Xe system apparently survived the NWA 5400 parent body processing in at least some mineral phases, although the exact iodine-carrier phase has yet to be identified.

Supported by NASA grant #NNX14A124G.

References: [1] Irving A. J. et al. (2009) 40th Lunar & Planet. Science Conf. A#2332. [2] Shukolyukov A. et al. (2010) 41th Lunar & Planet. Science Conf. A#1492. [3] Larouci N. et al. (2013) 76th Met. Soc. Meeting, A#5185. [4] Day J. M. D. et al. (2012) *Geochim. Cosmochim. Acta* 81, 94-128. [5] Burkhardt C. et al. (2015) 46th Lunar & Planet. Science Conf. #2732. [6] Amelin Y. and Irving A. J. (2011) 74th Met. Soc. Meeting, A#5197. [7] Shukolyukov A. et al. (2010) 41th Lunar & Planet. Science Conf. #1492. [8] Gilmour J. D. et al. (2009) *Meteoritics & Planet. Science* 44:573-579. [9] Bouvier A. and Wadhwa M. (2010) *Nature Geoscience* 3, 637-641.