

Gravity aspect reveals more differentiated southern hemisphere than northern hemisphere of Mars.

Gunther Kletetschka^{1,2}, Jaroslav Klokočník³, Jan Kostelecký^{4,5}, and Aleš Bezděk^{3,6},

¹Geophysical Institute, University of Alaska, Fairbanks, 903 N Koyukuk Drive, Fairbanks, AK, USA, gkletetschka@alaska.edu,

²Institute of Hydrogeology, Engineering Geology and Applied Geophysics, Faculty of Science, Charles University, Albertov 6, 12000 Prague 2, Czech Republic,

³Astronomical Institute, Czech Academy of Sciences, CZ 251 65 Ondřejov, Fričova 298, Czech Republic,

⁴Research Institute of Geodesy, Topography and Cartography, CZ 250 66 Zdiby 98, Czech Republic,

⁵ Faculty of Mining and Geology, VSB-TU Ostrava, CZ 708 33 Ostrava, Czech Republic,

⁶ Faculty of Civil Engineering, Czech Technical University in Prague, CZ 166 29 Praha 6, Czech Republic)



Abstract #6021

Introduction: We use the gravity aspects (descriptors) to study features on Mars. They are derived from the gravity field models of Mars (sets of spherical harmonic expansion of disturbing gravitational potential in terms of harmonic coefficients, also known as Stokes parameters) [1].

Results: Gravity aspect related to second derivative of the gravity potential (Fig. 1) revealed a significant difference in the crustal density distribution near the poles of Mars. Specifically, while the north and south polar crust region have similar distribution of gravity potential difference (-250 mGals to 350 mGals), related to first derivative of gravity potential (Δg , see [1] for exact definitions and Fig. 2), the northern region has significantly increased second derivative of the potential (T_{zz} see [1] for exact definitions), compared to southern crust polar region (Fig. 1).




Fig. 1. Fig. 1: The second derivative of the gravity potential T_{zz} [Etwos] in the polar regions of Mars. Plots are based on the global gravity field model JGMRO_120F [2]. Countours are from the MOLA MGS topography (black contours in [m] above reference surface).

Discussion: Northern and southern crust of Mars relates to Mars dichotomy. Northern crust has significantly lower elevation compared to southern crust of Mars. Our data indicate that the northern crust have near surface bodies with density contrast as much as 3 times larger than near surface crust in the southern polar region. Larger density contrast is consistent with the northern hemisphere being less differentiated, with contrasting densities, than the southern hemisphere with more uniform densities.




Fig. 2. Gravity aspects Δg for both northern and southern hemispheres. Here Δg [mGal], is plotted along with the MOLA topography.

Acknowledgments: GK was partially supported from the Czech Science Foundation 20-08294S, Ministry of Education, Youth and Sports LTAUSA 19141. Kurosh Karimi for imaging of VD of Mars.

References: [1] Klokočník J. et al. (2020), ISBN (10): 1-5275-4948-8; ISBN (13): 978-1-5275-4948-7. [2] Konopliv A. et al. (2020) Geophys. Res. Letts, 47.. doi: 10.1029/2020GL090568.

